

12

EUROPEAN PATENT APPLICATION

21 Application number: 89304620.1

51 Int. Cl.4: **C11D 3/39 , C11D 3/386 ,
C12N 15/00**

22 Date of filing: 08.05.89

30 Priority: 09.05.88 GB 8810954

43 Date of publication of application:
15.11.89 Bulletin 89/46

64 Designated Contracting States:
CH DE ES FR GB IT LI NL SE

71 Applicant: **UNILEVER PLC**
Unilever House Blackfriars
London EC4P 4BQ(GB)

84 GB

Applicant: **UNILEVER NV**
Burgemeester s'Jacobplein 1 P.O. Box 760
NL-3000 DK Rotterdam(NL)

84 CH DE ES FR IT LI NL SE

72 Inventor: Lagerwaard, Cornelis Abraham
Liguster 5
NL-3224 TM Hellevoetsluis(NL)
Inventor: Cornelissen, Johannes Mattheus
Platanendreef 27
NL-3137 CN Vlaardingen(NL)
Inventor: Swarthoff, Ton
Debussystraat 3
NL-3223 TE Hellevoetsluis(NL)
Inventor: Klugkist, Jan
Staverenhoeve 12
NL-3137 GN Vlaardingen(NL)
Inventor: Thom, David
Cherry Bank Manorial Road
Parkgate Wirral Merseyside(GB)

74 Representative: Ford, Michael Frederick et al
MEWBURN ELLIS & CO. 2/3 Cursitor Street
London EC4A 1BQ(GB)

EP 0 341 947 A1

54 Enzymatic detergent and bleaching composition.

57 An enzymatic detergent and bleaching composition contains a bleaching system which is stronger than the sodium perborate/TAED system. In such a composition a lipase is employed which is produced by cloning, by rDNA technique, the gene which encodes the lipase produced by the fungus Humicola lanuginosa and expressing that gene in Aspergillus oryzae as host. The lipase is used in an amount giving a lipolytic enzyme activity of from 100 to 0.005 lipolytic units/mg of the composition.

ENZYMATIC DETERGENT AND BLEACHING COMPOSITION

The present invention relates to an enzymatic detergent and bleaching composition comprising as essential ingredients a lipolytic enzyme and a bleaching system.

Detergent compositions comprising lipase are known. For example GB 1 372 034 (Unilever) discloses lipase from *Pseudomonas* in specific nonionic-containing detergent compositions for soaking fabrics.

5 USP 3 950 277 (Procter & Gamble) also describes fabric-soaking compositions: the described compositions comprise lipase and lipase activators and a number of lipases from microorganism and other sources are mentioned: those particularly mentioned as preferred are Amano CE, Amano M-AP, Takeda 1969-4-9, and Meito MY-30 lipases, but no indications are given of the form in which the lipase is to be prepared or used.

10 USP 4 011 169/NL 74 08763 (Procter & Gamble) describes the use of a similar range of enzymes in the preparation of additives for washing agents (detergent compositions).

Examples of known lipase-containing detergent compositions are provided by EP 0 205 208 and 0 208 390 (Unilever), which relate to lipases related to those from *Ps. fluorescens*, *P. gladioli* and *Chromobacter* in detergent compositions.

15 EP 0 214 761 (Novo) and EP 0 258 068 (Novo), each give detailed description of lipases from certain microorganisms, and also give certain uses in detergent additives and detergent compositions for the enzymes described. EP 0 214 761 gives detailed description of lipases derived from organisms of the species *Pseudomonas cepacia*, and certain uses therefor. EP 0 258 068 gives detailed description of lipases derived from organisms of the genus *Thermomyces/Humicola*, and certain uses therefor.

20 Also believed to be in use in certain areas is a lipase-containing granular detergent composition containing about 37% detergent actives including 5% nonionic detergent and the remainder substantially anionic detergent, about 16% zeolite, about 60 LU/g lipase, plus protease and other normal detergent additives.

Further examples of known lipase-containing detergent compositions are provided by JA 63-078000 25 (1988) (Lion Corp/K Mukoyama et al) which discloses properties and uses of a *Pseudomonas* lipase, including use in a lipase-containing system based on 10-40 % surfactant (e.g., sodium C14-C18 alpha-olefin sulphate), as well as other conventional detergent ingredients.

In EP 0 268 456 (Clorox), there is described in connexion with Table 10(b) an experimental washing solution containing lipase and about 1 microgram/ml sodium dodecyl sulphate.

30 In US Patent No 4,707,291 (hereby incorporated by reference herein), detergent compositions have been described which contain a special class of lipolytic enzymes. These compositions may also contain a bleaching agent such as sodium perborate, either as such or in admixture with a low temperature bleach activator, e.g., tetraacetyl-ethylene-diamine (TAED). The lipases mentioned in US Patent No 4,707,291 are significantly less affected by a bleaching system than other lipases. These bleaching systems comprise 35 sodium perborate and TAED.

We have now surprisingly found that a certain type of lipase, defined herein below, is quite compatible with bleaching systems which are stronger than the sodium perborate/TAED system, such systems being defined in more detail hereafter. This type of lipase is more stable in the presence of these bleaching systems than some of the lipases mentioned US Patent No 4,707,291 and is in particular more stable than 40 the lipases mentioned in US Patent No 4,707,291 in the presence of proteases in these bleaching systems.

The lipase used in the present invention is a lipase produced by cloning, by rDNA technique, the gene encoding the lipase produced by the fungus *Humicola lanuginosa* and expressing the gene in *Aspergillus oryzae* as host. This product is manufactured and commercially available from Novo Industri A/S, Denmark, under the trade name Lipolase (see 'Biotechnology Newswatch', published 7th March 1988, page 6). In the 45 relevant manufacturer's technical leaflet on Lipolase (Trade Mark) it is said that this lipase is not stable in the presence of e.g., hypochlorite as bleaching agent.

Equivalence of a lipase enzyme produced by rDNA technique to the lipase defined above can be determined in practice for example by an immunological reaction of identity between the two lipase materials and/or by determination of substantial aminoacid sequence homology therebetween. In some such 50 equivalent lipases, of which the use is within the scope of the invention, the pattern of glycosylation of the lipase protein differs.

EP 0 305 216 (Novo) also describes suitable lipase enzymes for use in the practice of this invention, and is hereby incorporated by reference herein.

The lipase used in the present invention is included in the detergent and bleaching composition in such an amount that the final composition has a lipolytic enzyme activity of from 100 to 0.005 LU/mg, preferably

25 to 0.05 LU/mg of the composition.

A Lipase Unit (LU) is that amount of lipase which produces 1/micromol of titratable fatty acid per minute in a pH stat. under the following conditions: temperature 30 °C; pH = 9.0; substrate is an emulsion of 3.3 wt.% of olive oil and 3.3% gum arabic, in the presence of 13 mmol/l Ca⁺⁺; and 20 mmol/l NaCl in 5 mmol/l Tris-Buffer. An alternative definition of the lipase unit is given in EP 0 258 068 (Novo), which mentions (inter alia) lipases suitable for use in the practice of this invention and is hereby incorporated by reference.

Naturally, mixtures of the above lipase with other known lipases can be used. The lipase(s) can be used in its (their) non-purified form or in a purified form, e.g., purified with the aid of well-known adsorption methods, such as phenyl sepharose adsorption techniques.

The bleaching system used according to the present invention is stronger than the sodium perborate/TAED system. This latter system, through a perhydrolysis reaction, forms a peroxyacid, i.e., peracetic acid, but at a rather low rate. The bleaching systems according to the present invention must be stronger than this sodium perborate/TAED system, by which is to be understood that the system either is based on a peracid (inorganic or organic) which is stronger than the peracetic acid or yields, on perhydrolysis, an organic peracid, including peracetic acid, faster than the sodium perborate/TAED system. The bleaching system may consist of a bleaching agent as such or may consist of a bleaching agent together with a bleach precursor. As bleaching agent as such alkali metal monopersulphates, furthermore organic peracids such as diperoxy dodecanedioic acid, diperoxy tetradecanedioic acid, diperoxyhexadecanedioic acid, mono- and diperazelaic acid, mono- and diperbrassylic acid, monoperoxy phthalic acid, perbenzoic acid, can be used, either as acid or in the form of their salts.

When a system comprising a bleach precursor is used, this system comprises a bleaching agent which reacts with a bleach precursor to form a peracid in solution faster than the sodium perborate/TAED system. By faster is meant that the precursor will have a rate of peroxy acid release of at least 2 (two) times, preferably at least 5 (five) times faster than TAED under the same conditions.

Typical examples of such systems are sodium perborate with sodium nonanoyloxy benzene sulphonate or sodium trimethyl hexanoyloxy benzene sulphonate or sodium acetoxy benzene sulphonate or sodium benzoyloxy benzene sulphonate.

The preferred systems of the present invention are sodium perborate with sodium nonanoyloxy benzene sulphonate, diperoxy dodecane dioic acid or monopersulphate.

In general, the amount of the bleaching system in the composition varies from 1-50%, usually from 5-40% by weight. When a bleach precursor is present, the molar ratio of the bleach precursor to the percompound such as sodium perborate varies from 1:1 to 1:35, preferably from 1:2 to 1:20. Mixtures of various bleaching agents and various bleach precursors in accordance with the invention can also be used.

The compositions of the present invention may furthermore contain one or more detergent active materials, such as soaps, anionic, nonionic, cationic and zwitterionic synthetic detergents or mixtures thereof. Usually the amount of detergent active material present in the composition will range from 1-50%, preferably 2-40% and particularly preferably 5-30% by weight. Suitable examples of detergent active materials can be found in Schwartz, Perry and Berch "Surface Active Agents and Detergents", Vol I (1949) and Vol II (1958) and M Schick "Nonionic Surfactants" Vol I (1967).

The compositions may furthermore include the usual detergent ingredients in the usual amounts. They may be unbuilt or built, and may be of the zero-P type (i.e., not containing phosphorus-containing builders). Thus, the compositions may contain from 1-60%, preferably from 5-30% by weight of one or more organic and/or inorganic builders. Typical examples of such builders are the alkali metal ortho-, pyro- and tripolyphosphates, alkali metal carbonates, either alone or in admixture with calcite, alkali metal citrates, alkali metal nitrilotriacetates, carboxymethyloxy succinates, zeolites, polyacetal carboxylates and so on.

The compositions may furthermore comprise lather boosters, foam depressors, anti-corrosion agents, soil-suspending agents, sequestering agents, anti-soil redeposition agents, perfumes, dyes, stabilizing agents for the enzymes and bleaching agents and so on. They may also comprise enzymes other than lipases, such as proteases, amylases, oxidases and cellulases. In this respect it has been found that, whereas proteases, are often affected by strong bleaches, in the present invention, when used together with the lipases of the present invention, the overall performance of the enzyme system is often not significantly affected.

In general, the compositions may comprise such other enzymes in an amount of 0.01-10% by weight. For proteases, the amount, expressed in proteolytic activity, is usually from 0.1-50 GU/mg based on the final composition.

A GU is a glycine unit, which is the amount of proteolytic enzyme which under standard incubation conditions produces an amount of terminal NH₂-groups equivalent to 1 microgram/ml of glycine.

The compositions of the present invention can be formulated in any desired form, such as powders, bars, pastes, liquids, etc.

The invention will further be illustrated by way of Example. The invention described herein extends to modifications and variations as will be apparent to the skilled reader, and combinations and subcombinations of the features mentioned herein.

EXAMPLE 1

10

The stability of various lipases in the presence of a bleaching system was measured as follows:

To a solution of 4 g/l of a detergent composition* and 0.03 g/l Dequest 2041 in water with a hardness of 30 ° FH and a temperature of 30 ° C, an amount of lipase is added to obtain 15-20 lipase units/ml.

The pH is adjusted with NaOH to pH 10.0 at 30°C. At t=0 a bleach system is added:

15

- a) 292 mg/l TAED (65% pure) and 700 mg/l sodium perborate monohydrate or
- b) 1880 mg/l DPDA (12% pure) or
- c) 822 mg/l SNOBS (80% pure) and 1500 mg/l sodium perborate monohydrate or
- d) 506 mg/l MPS (in the form of the commercial product Caroate R) or
- e) 475 mg/l P15 (95% pure) and 700 mg/l sodium perborate monohydrate.

20

This yields 1.5 mmolar peracid in solution for all bleach systems. The lipase stability is measured by determining the residual lipase activity with the pH-stat. method.

25

30

Dequest 2041	= ethylene diamine tetra (methylene phosphonic acid)
TAED	= tetraacetyl ethylene diamine
DPDA	= diperoxy dodecanedioic acid
SNOBS	= sodium nonanoyloxy benzene sulphonate
MPS	= sodium monopersulphate
P15	= sodium benzoyloxy benzene sulphonate

35

40

45

50

55

* The detergent composition had the following formulation:

	% by weight
Sodium dodecyl benzene sulphonate	6.5
C14-C15 primary alcohol, condensed with 11 moles of ethylene oxide	2.0
Sodium stearate	1.0
Sodium silicate	7.0
Sodium carboxymethyl cellulose	0.5
Na ₂ S ₂ O ₄	37.0
Pentasodium triphosphate	15.0
Trisodium orthophosphate	5.0
Fluorescer	0.2
Ethylene diamine tetraacetic acid	0.5
Water	6.2
Dyes	0.01

The following results were obtained :																			
Lipase ex.	Trade-name	No bleach activity			TAED/perb. activity			SNOBS/perb. activity			DPDA activity			MPS activity			P15 activity		
		10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)
Humicola lanuginosa	Lipolase	100	99	>30	100	85	>30	95	83	>30	95	70	>30	94	72	>30	100	96	>30
Pseudomonas gladioli	-	97	97	>30	97	92	>30	90	80	>30	100	99	>30	89	80	>30	92	77	>30
Chromobacter viscosum	Toyo Jozo	100	95	>30	100	86	>30	80	58	>30	107	98	>30	85	62	>30	93	63	>30
Pseudomonas fluorescens	Amano P	100	98	>30	92	100	>30	85	80	>30	109	100	>30	89	85	>30	102	83	>30
Pseudomonas cepacia	ex NOVO	93	90	>30	98	94	>30	89	81	>30	100	98	>30	97	96	>30	95	84	>30
Aspergillus niger	Amano AP 6	110	90	>30	100	64	43	14	<5	2	62	<5	12	78	40	24	30	9	7
Mucor Miehei	SP 225	67	46	28	95	58	35	81	46	28	86	68	>30	13	5	4	73	44	26

residual lipase activity (% of input)
t1/2 = half life time

EXAMPLE 2

Example 1 was repeated, but now in the presence of 20 GU (glycine unit)/ Savinase , a proteolytic enzyme ex NOVO. The following results were obtained :																		
Lipase	No bleach activity			TAED/perb. activity			SNOBS/perb. activity			DPDA activity			MPS activity			P15 activity		
	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)	10 min	30 min	t1/2 (min)
Lipolase	100	100	>30	100	88	>30	94	85	>30	96	90	>30	94	72	>30	100	96	>30
Ps. gladioli	81	56	37	76	51	31	76	55	36	90	70	>60	63	47	20	81	42	26
Amano P	51	17	10	56	20	12	40	16	6	60	24	12	43	27	8	55	15	11
Ps. cepacia	65	35	18	72	38	19	65	34	19	59	42	18	54	32	12	65	28	17
Toyo Jozo	79	48	30	71	38	18	72	47	28	82	52	33	38	22	8	74	29	17
Amano AP6	96	83	>30	82	38	25	<5	<5	3	61	15	12	91	79	>30	55	24	11
Esterase MM	64	21	13	38	15	8	43	12	8	68	25	16	10	<5	5	74	25	17

" in an experiment with Kazusase, a proteolytic enzyme ex Showa Denko, the following results were obtained: 95% 84% >30.

" with Alcalase instead of Savinase, the results were as follows: 97% 93% >30.

" with Alcalase instead of Savinase, the results were as follows: 90% 76% >30.

" with Kazusase instead of Savinase, the results were as follows: 99% 89% >30.

residual lipase activity (% of input)

t/2 = half life time

Claims

- 5 1. A detergent composition comprising from 1-50% by weight of one or more detergent-active materials, from 0-60% by weight of a builder, from 1-50% by weight of a bleaching agent and lipolytic enzyme in an amount of 0.005-100 lipolytic units per milligram of the composition, characterised in that the bleaching agent is based on an inorganic or organic peracid or salt thereof which is stronger than peracetic acid or comprises a bleaching agent and a bleach precursor which yields, on perhydrolysis, a peracid faster
 10 than the system comprising sodium perborate + tetraacetyl ethylene diamine, and the lipolytic enzyme is a lipase produced by cloning by rDNA technique the gene encoding the lipase produced by Humicola lanuginosa (syn. Thermomyces lanuginosus) and expressing the gene in Aspergillus oryzae as host.
2. A composition according to Claim 1, characterised in that the bleaching agent is an alkali metal persulphate.
- 15 3. A composition according to Claim 1 characterised in that the bleaching agent is selected from the group consisting of diperoxy dodecanedioic acid, diperoxy tetradecanedioic acid, diperoxyhexadecane diolc acid, mono- and diperazelaic acid, mono- and diperbrassylic acid, monoperoxy phthalic acid, perbenzoic acid, and their salts.
4. A composition according to Claim 1, characterized in that the bleaching agent comprises a bleaching
 20 agent and a bleach precursor which forms a peracid in solution at least two times faster than tetraacetyl ethylene diamine under the same conditions.
5. A composition according to Claim 4, characterised in that the bleaching agent comprises sodium perborate and a bleach precursor selected from the group consisting of sodium nonanoyloxy benzene sulphonate, sodium trimethyl hexanoyloxy benzene sulphonate, sodium acetoxo benzene sulphonate and
 25 sodium benzoyloxy benzene sulphonate.
6. A composition according to any of Claims 1-5, characterized in that it further contains a proteolytic enzyme in an amount of 0.1-50 GU/mg of the composition.
7. A composition according to any of Claims 1-6, characterised in that the lipase enzyme is Lipolase (Trade Mark).

30

35

40

45

50

55



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
P, A	EP-A-0 271 152 (UNILEVER) * Whole document *	1	C 11 D 3/39 C 11 D 3/386 C 12 N 15/00
A	LU-A- 65 030 (PROCTER & GAMBLE) * Claims; pages 10,11 *	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 11 D C 12 N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 01-08-1989	Examiner PFANNENSTEIN H. F.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	